

Attorney's Docket: 1999DE132
Serial No.: 09/722,760
Group: 1756

AMENDMENTS TO THE CLAIMS

1. (Currently Amended) A method of imparting, controlling or improving the charge of an electrophotographic toner or developer, or an electret material, comprising the step of adding as a charge control agent a structured silicate salt wherein the cation is a low molecular weight, nonpolymeric organic cation or a combination of a low molecular weight, nonpolymeric organic cation with NH_4^+ , H_3O^+ , an alkali metal, an alkaline earth metal, an earth metal or with a transition metal and the anion is an island, cyclic, group, chain, ribbon, laminar or matrix silicate or a combination thereof to a binder of an electrophotographic toner or developer or of an electret material.

2) (Cancelled)









3) (Cancelled)

4) (Previously Presented) The method as claimed in claim 1, wherein the low molecular weight organic cation is a substituted, phosphonium, thionium or triphenylcarbonium ion or a cationic metal complex.



5) (Currently Amended) The method as claimed in claim 1, wherein the low molecular weight organic cation is an ammonium ion having one of the formulae (a) - (j)

BEST AVAILABLE COPY

Attorney's Docket: 1999DE132
Serial No.: 09/722,760
Group: 1756




Attorney's Docket: 1999DE132
 Serial No.: 09/722,760
 Group: 1756



In which

R^1 to R^{18} are identical or different and represent hydrogen, CN, $(\text{CH}_2)_{1-18}\text{CN}$, halogen, branched or unbranched C_1 - C_{32} -alkyl, mono- or polyunsaturated C_2 - C_{32} -alkenyl, C_1 - C_{22} -alkoxy, C_1 - C_{22} -hydroxyalkyl, C_1 - C_{22} -halogenoalkyl, C_2 - C_{22} -halogenoalkenyl, C_1 - C_{22} -aminoalkyl, $(\text{C}_1$ - C_{12})-trialkyl-ammonium- $(\text{C}_1$ - C_{22})-alkyl; $(\text{C}_1$ - C_{22})-alkylene- $(\text{C}=\text{O})\text{O}$ - $(\text{C}_1$ - C_{32})-alkyl, $(\text{C}_1$ - C_{22})-alkylene- $(\text{C}=\text{O})\text{O}$ -aryl, $(\text{C}_1$ - C_{22})-alkylene- $(\text{C}=\text{O})\text{NH}$ - $(\text{C}_1$ - C_{32})-alkyl, $(\text{C}_1$ - C_{22})-alkylene- $(\text{C}=\text{O})\text{NH}$ -aryl, $(\text{C}_1$ - C_{22})-alkylene- $\text{O}(\text{CO})$ - $(\text{C}_1$ - C_{32})-alkyl, $(\text{C}_1$ - C_{22})-alkylene- $\text{O}(\text{CO})$ -aryl, $(\text{C}_1$ - C_{22})-alkylene- $\text{NH}(\text{C}=\text{O})$ - $(\text{C}_1$ - C_{32})-alkyl, or $(\text{C}_1$ - C_{22})-alkylene- NHCO -aryl,

wherein






are optionally inserted into the acid ester or acid amide bonds;

$[(\text{C}_1$ - C_{12})-alkylene- $\text{O}]_{1-100}$ -H; aryl, $(\text{C}_1$ - C_{18})-alkylenearyl; $-(\text{O}-\text{SiR}'_2)_{1-32}-\text{O}-\text{SiR}'_3$, in which R' has the meaning C_1 - C_{12} -alkyl, phenyl, benzyl or C_1 - C_{12} -alkoxy; heterocyclyl, or C_1 - C_{18} -alkylene-heterocyclyl, wherein the aryl and heterocyclyl radicals are optionally mono- or polysubstituted on carbon atoms or heteroatoms by C_1 - C_{12} -alkyl, C_1 - C_4 -alkenyl, C_1 - C_4 -alkoxy, hydroxy- $(\text{C}_1$ - C_4)-alkyl, amino- $(\text{C}_1$ - C_4)-alkyl, C_1 - C_4 -alkylimino, carboxyl, hydroxyl, amino, nitro, cyano, halogen, C_1 - C_{12} -acyl, C_1 - C_4 -halogenoalkyl, C_1 - C_4 -alkylcarbonyl, C_1 - C_4 -alkylcarbonyloxy, C_1 - C_4 -alkoxycarbonyl, C_1 - C_4 -alkylaminocarbonyl, C_1 - C_4 -alkylcarbonylimino, C_6 - C_{10} -arylcarbonyl, aminocarbonyl, aminosulfonyl, C_1 - C_4 -alkylaminosulfonyl, phenyl, naphthyl, or heteroaryl;

Attorney's Docket: 1999DE132
 Serial No.: 09/722,760
 Group: 1756

R^{19} represents C_4 - C_{11} -alkylene, $-(C_2H_4-O-)_{1-17}-(CH_2)_{1-2}$, $-(C_2H_4-NR-)_{1-17}$ or $-(CH_2)_{1-2}$,
 in which R is hydrogen or C_1 - C_{12} -alkyl;
 X has the meaning of Y, or $-\text{CO}-\text{CH}_2-\text{CO}-$,



Y has the meaning $\begin{array}{c} \text{C} \\ \parallel \\ \text{O} \end{array}$, $\begin{array}{c} \text{C} \\ \parallel \\ \text{S} \end{array}$, $\begin{array}{c} \text{C} \\ \parallel \\ \text{NH} \end{array}$, $-(\text{CH}_2)_{1-18}$,

Attorney's Docket: 1999DE132
 Serial No.: 09/722,760
 Group: 1756



or o-, p-, m-(C₆-C₁₄)-arylene or (C₄-C₁₄)-heteroarylene with 1, 2, 3 or 4 heteroatoms selected from the group consisting of N, O, S and a combination thereof;

R⁶⁰ represents C₁-C₃₂-acyl, C₁-C₂₂-alkyl, C₂-C₂₂-alkenyl, C₁-C₁₈-alkylene-C₆-C₁₀-aryl, C₁-C₂₂-alkylene-heterocyclyl, C₆-C₁₀-aryl or (C₄-C₁₄)-heteroaryl with 1, 2, 3 or 4 heteroatoms selected from the group consisting of N, O, S, and a combination thereof;

R⁶¹ and R⁶⁴ represent -(CH₂)₁₋₁₈-, C₁-C₁₂-alkylene-C₆-C₁₀-arylene, C₆-C₁₀-arylene, or C₆-C₁₂-alkylene-heterocyclyl;


Z represents -NH- or -O-;

A₁⁻ and A₃⁻ represent -COO⁻, -SO₃⁻, -OSO₃⁻, -SO₂⁻, -COS⁻ or -CS₂⁻;

A₂ represents -SO₂Na, -SO₃Na, -SO₂H, -SO₃H or hydrogen;

R⁶⁹ and R⁷⁰ independently of one another represent hydrogen, C₁-C₃₂-alkyl, in which the alkyl chain optionally contain one or more of the groups selected from the group consisting of -NH-CO-, -CO-NH-, -CO-O-, or -O-CO-; C₁-C₁₈-alkylene-aryl, C₆-C₁₈-alkylene-heterocyclyl, C₁-C₁₈-hydroxyalkyl, C₁-C₁₈-halogenoalkyl, aryl, -(CH₂)₃-SO₃⁻,
or

Attorney's Docket: 1999DE132
 Serial No.: 09/722,760
 Group: 1756



R⁷¹ and R⁷² represent $-(\text{CH}_2)_{1-12}-$; and
 R⁷³ and R⁷⁴ represent hydrogen or C₁-C₂₂-alkyl.

6) (Original) The method as claimed in claim 5, wherein R¹ to R¹⁸ denote hydrogen, CN, CH₂-CN, CF₃, C₁-C₂₂-alkyl, C₂-C₁₈-alkenyl, C₁-C₁₈-alkoxy, C₁-C₁₈-hydroxy-alkyl, C₁-C₁₈-halogenoalkyl, C₂-C₁₈-halogenoalkenyl, C₁-C₁₈-aminoalkyl, (C₁-C₆)-trialkylammonium-(C₁-C₁₈)-alkyl, (C₁-C₁₈)-alkylene-O(C=O)-(C₁-C₂₂)-alkyl, (C₁-C₁₈)-alkylene-O(C=O)-phenyl, (C₁-C₁₈)-alkylene-NHCO-(C₁-C₂₂)-alkyl, (C₁-C₁₈)-alkylene-NHCO-phenyl, (C₁-C₁₈)-alkylene-(C=O)O-(C₁-C₂₂)-alkyl, (C₁-C₁₈)-alkylene-(C=O)O-phenyl, (C₁-C₁₈)-alkylene-(C=O)NH-(C₁-C₂₂)-alkyl, (C₁-C₁₈)-alkylene-CONH-phenyl, benzyl, phenyl, naphthyl, C₁-C₁₂-alkylene-heterocyclyl;
 R¹⁹ denotes C₄-C₅-alkylene, $-(\text{C}_2\text{H}_4\text{-O})_{1-6}-$, $-(\text{CH}_2)_{1-2}-$ or $-(\text{C}_2\text{H}_4\text{-NH})_{1-6}-$, $-(\text{CH}_2)_{1-2}-$;
 R⁶⁰ denotes C₁-C₁₈-acyl, C₁-C₁₈-alkyl, C₂-C₁₈-alkenyl, C₁-C₁₂-alkylene-phenyl, C₁-C₁₈-alkylene-pyridyl, phenyl or pyridyl;
 R⁶¹ and R⁶⁴ denote $-(\text{CH}_2)_{1-12}-$, C₁-C₈-alkylene-phenylene, phenylene or C₁-C₈-alkylenepyridylene or piperidylene;
 R⁷¹ and R⁷² denote $-(\text{CH}_2)_{1-8}-$ and
 R⁷³ and R⁷⁴ denote hydrogen or (C₁-C₁₈)-alkyl.

7) (Previously Presented) The method as claimed in claim 1, wherein the low molecular weight organic cation is an ammonium ion which is an aliphatic or

Attorney's Docket: 1999DE132
Serial No.: 09/722,760
Group: 1756

aromatic 5- to 12-membered heterocyclic radical with 1 to 4 atoms selected from the group consisting of N, O and S, or a combination thereof, belonging to the rings.

8) (Original) The method as claimed in claim 7, wherein the heterocyclic radical is pyridinium, pyridazinium, pyrimidinium, pyrazinium, purinium, tetraazaporphyrinium, piperidinium, morpholinium, tetrazonium, triaza-cyclononanium or tetraaza-cyclododecanium.

9) (Original) The method as claimed in claim 4, wherein the cationic metal complex is a metal carboxylate, metal salicylate, metal sulfonate, 1:1 metal-azo complex or a metal dithiocarbamate.

10) (Previously Presented) The method as claimed in claim 9, wherein the metal is selected from the group consisting of Al, Mg, Ca, Sr, Ba, TiO, VO, Cr, V, Ti, Zr, Sc, Mn, Fe, Co, Ni, Cu, Zn and ZrO.

11) through 16) (Cancelled)

17) (Previously Presented) The method as claimed in claim 9, wherein the metal is selected from the group consisting of Al, Mg, Ca, Sr, Ba, TiO, VO, Cr, V, Ti, Zr, Sc, Mn, Fe, Co, Ni, Cu, Zn and ZrO, and the metal complex contains one or more further ligands.

18) through 21) (Cancelled)

22) (Previously Presented) A method of imparting, controlling or improving the charge of an electrophotographic toner or developer, or an electret material comprising the step of adding a charge control agent to a binder of an

Attorney's Docket: 1999DE132
Serial No.: 09/722,760
Group: 1756

electrophotographic toner or developer or of an electret material, wherein the charge control agent is distearyl dimethyl ammonium bentonite.

23) (Previously Presented) A method of imparting, controlling or improving the charge of an electrophotographic toner or developer, or an electret material, comprising the step of adding as a charge control agent a structured silicate salt wherein the cation is a low molecular weight organic cation or a combination of a low molecular weight organic cation with NH_4^+ , H_3O^+ , an alkali metal, an alkaline earth metal, an earth metal or with a transition metal and the anion is an island, cyclic, group, chain, ribbon, laminar or matrix silicate or a combination thereof to a binder of an electrophotographic toner or developer or of an electret material, wherein the charge control agent imparts either a positive or negative charge.

24. (New) The method as claimed in claim 7, where 2 to 8 rings are fused:

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ ~~LINES OR MARKS ON ORIGINAL DOCUMENT~~
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.